



For questions or emergency care, please contact us at 985-327-5905.

POSTOPERATIVE INSTRUCTIONS

Ventilation (PE) Tubes - Pediatric

A ventilation (PE) tube is placed through a small incision in the eardrum to allow better aeration of the middle ear space. This is usually done to treat recurrent ear infections and/or fluid in the middle ear.

What to Expect:

- There is usually an initial fussy period of approximately 30 minutes following placement of tubes. Much of this is due to the initial confusion and disorientation of waking up after anesthesia. This should quickly pass and is commonly followed by a sleepy period. Your child should return to a normal routine later in the day but may continue to have periods of irritability.
- Drainage from the ears may occur for a few days after surgery especially with the use of antibiotic ear drops. It may appear clear, pink, or blood-tinged. At some point during the time your child has tubes, you may see additional drainage of fluid from the ears. This is most common during a viral illness. Antibiotic ear drops may be used alone to help clear this drainage.
- You may choose to place a dry cotton ball in the outer ear to absorb the drainage.
- Ear plugs may or may not be recommended after surgery. Consult with your surgeon regarding ear plug use.

Medication:

- After surgery, you may or may not need to use antibiotic drops in your child's ear. If drops are used, please follow the recommendation on your prescription.
- If needed, you may administer acetaminophen (Tylenol) or ibuprofen (Motrin) products as directed on the package instructions.

When to Call the Doctor:

- If your child develops a fever over 101°.
- If there is a steady trickle of blood or if drainage continues beyond 1 week despite use of antibiotic ear drops.
- If you need a refill prescription of antibiotic ear drops called to your pharmacy.

For questions or emergency care, please contact us at 985-327-5905.